

New Realisations of W Algebras and W Strings

H. Lu,^{*} C.N. Pope,^{*} S. Schrans[◇] and X.J. Wang

*Center for Theoretical Physics, Texas A&M University,
 College Station, TX 77843-4242, USA.*

ABSTRACT

We discuss new realisations of W algebras in which the currents are expressed in terms of two arbitrary commuting energy-momentum tensors together with a set of free scalar fields. This contrasts with the previously-known realisations, which involve only one energy-momentum tensor. Since realisations of non-linear algebras are not easy to come by, the fact that this new class exists is of intrinsic interest. We use these new realisations to build the corresponding W -string theories and show that they are effectively described by two independent ordinary Virasoro-like strings.

Available from hep-th@xxx/9203024

^{*} Supported in part by the U.S. Department of Energy, under grant DE-FG05-91ER40633.

[◇] Onderzoeker I.I.K.W.; On leave of absence from the Instituut voor Theoretische Fysica, K.U. Leuven, Belgium.

1. Introduction

The quantisation of two-dimensional matter systems with local W symmetries gives rise to new string theories, called W strings [1,2,3,4,5,6]. Traditional Virasoro string theory acquires its physical interpretation when the Virasoro symmetry is realised by a set of free scalar fields, which then describe the coordinates of the target spacetime. Similarly, the physical interpretation of a W string depends upon the realisation of the W symmetry. For the bosonic WA_n and WD_n strings based on the A_n and D_n classical simply-laced Lie algebras, the realisations that have been found so far are given in terms of an arbitrary energy-momentum tensor T^{eff} , together with a set of free scalar fields φ_i which are “frozen” by the physical-state conditions. If T^{eff} is itself realised in terms of free scalar fields X^μ , then these scalars acquire the interpretation of being target-spacetime coordinates. The physical spectrum of these W strings thus closely resembles the spectrum of the ordinary Virasoro string, but with a non-critical value of the central charge, and non-standard values for the spin-2 intercept [5,6].

In the first part of this paper we obtain new, inequivalent, realisations for the WA_n and WD_n algebras with $n \geq 3$. These realisations are given in terms of *two* arbitrary energy-momentum tensors T_+ and T_- (which commute with each other, and have the same central charge), together with $(n - 2)$ extra free scalar fields. They are the first examples of realisations that are not of the form described in the previous paragraph. Since the W algebras are non-linear, realisations are not easy to come by; in particular, unlike the Virasoro algebra, one cannot obtain new realisations simply by tensoring together old ones. Thus these new realisations are of intrinsic interest for their own sake.

In the second part of this paper we use these realisations to construct the corresponding W -string theories, and show that they are quite different from the W -string theories that have previously been constructed. The physical spectrum is effectively described by two Virasoro-type strings living in two independent Minkowski spacetimes.

2. Reduction

In order to explain how the new realisations of WA_n and WD_n arise, it is useful to begin by reviewing how the previously-known realisations with only one arbitrary energy-momentum tensor are constructed.* The starting point is the Miura transformation, which provides a realisation in terms of n scalar fields [7,8]. For WA_n , $(n - 1)$ of these scalars appear in the generating currents only *via* the Miura transformation of WA_{n-1} . Thus the currents of WA_n can be realised in terms of those of WA_{n-1} , together with the remaining scalar field. This remarkable property is a consequence of the detailed form of the Miura transformation. It was first noticed for $WA_2 = W_3$ in [9], and generalised to all WA_n in [2,5]. Applying this reduction recursively, one can therefore realise the currents of WA_n in terms of $(n - 1)$ scalar fields together with an arbitrary energy-momentum tensor T^{eff} which generates WA_1 , the Virasoro algebra. This reduction procedure can be summarised by the diagram

$$WA_n \rightarrow WA_{n-1} \rightarrow \cdots \rightarrow WA_2 \rightarrow WA_1 = \text{Virasoro}. \quad (2.1)$$

For WD_n , it was shown in [6] that a similar reduction is possible, with the currents being expressed in terms of those of WD_{n-1} together with an additional free scalar. Since $D_2 \cong A_1 \times A_1$ is not simple, one usually restricts the discussion of WD_n algebras to the cases $n \geq 3$ [8]. Thus WD_n can be realised in terms of the currents of WD_3 together with $(n - 3)$ free scalar fields. Since $D_3 \cong A_3$, and thus $WD_3 \cong WA_3$, we

* Recall that WA_n is the W_{n+1} algebra, generated by currents of spins $2, 3, \dots, n + 1$ [7]. The WD_n algebra is generated by currents of spins $2, 4, 6, \dots, 2n - 2$ and n [8].

may then continue the reduction procedure by following the route given in (2.1) to descend from WA_3 to WA_1 . This can be summarised by the diagram

$$WD_n \rightarrow WD_{n-1} \rightarrow \cdots \rightarrow WD_3 \cong WA_3 \rightarrow WA_2 \rightarrow WA_1 = \text{Virasoro}. \quad (2.2)$$

Thus WD_n can be realised in terms of an arbitrary energy-momentum tensor and $(n-1)$ additional free scalars.

Although one does not usually consider the WD_2 algebra, the Miura transformation for WD_n allows one to follow the reduction scheme that we have described above for one further step, to express the currents of WD_n in terms of those of WD_2 together with $(n-2)$ additional free scalars. Since $D_2 \cong A_1 \times A_1$, the algebra WD_2 is isomorphic to the direct product of two independent Virasoro algebras. Thus the currents of WD_n may be expressed in terms of $(n-2)$ free scalars together with two arbitrary commuting energy-momentum tensors T_+ and T_- . The detailed form of the Miura transformation imposes the condition that T_+ and T_- should have the same central charge. The reduction procedure can be summarised by the diagram

$$WD_n \rightarrow WD_{n-1} \rightarrow \cdots \rightarrow WD_3 \rightarrow WD_2 = \text{Virasoro} \times \text{Virasoro}. \quad (2.3)$$

Exploiting the isomorphism between D_3 and A_3 in the opposite direction to that which we used in (2.2), we may also construct an alternative reduction procedure for WA_n , given by

$$WA_n \rightarrow WA_{n-1} \rightarrow \cdots \rightarrow WA_3 \cong WD_3 \rightarrow WD_2 = \text{Virasoro} \times \text{Virasoro}. \quad (2.4)$$

Thus the currents of WA_n (with $n \geq 3$) may be expressed in terms of $(n-2)$ free scalars together with the two arbitrary energy-momentum tensors T_+ and T_- .

We shall now illustrate this reduction procedure in more detail. A realisation of WD_n in terms of n free scalar fields $(\varphi_1, \dots, \varphi_n)$ has been given by Fateev and Lukyanov [8]. The spin- n current is given by the Miura-type transformation

$$f^{(n)}(z) = (\alpha_0 \partial - \partial \varphi_n) (\alpha_0 \partial - \partial \varphi_{n-1}) \cdots (\alpha_0 \partial - \partial \varphi_2) \partial \varphi_1, \quad (2.5)$$

where, as usual, normal ordering is assumed. The remaining currents $W_{2k}^{(n)}$ of spin $2k$, with $k = 1, 2, \dots, n-1$, can then be read off from the poles of order $(2n-2k)$ in the operator-product expansion $f^{(n)}(z)f^{(n)}(w)$ [8]. The reduction procedure $WD_n \rightarrow WD_{n-1}$ can now easily be understood by observing from (2.5) that we may rewrite $f^{(n)}$ as

$$f^{(n)} = \alpha_0 \partial f^{(n-1)} - \partial \varphi_n f^{(n-1)}, \quad (2.6)$$

where $f^{(n-1)}$ is the spin- $(n-1)$ current of WD_{n-1} . It then follows that the $W_{2k}^{(n)}$ currents can, likewise, be written in terms of the WD_{n-1} currents together with the extra scalar field φ_n . Applying this reduction procedure recursively one may then realise the WD_n algebra in terms of the currents generating WD_2 together with the $(n-2)$ free scalars $(\varphi_3, \dots, \varphi_n)$. As explained previously, WA_n can also be realised in terms of the currents generating WD_2 together with $(n-2)$ free scalars, by following the route given in (2.4). Since the key step in these reduction schemes for both WA_n and WD_n is the reduction from WD_3 (which is isomorphic to WA_3) to WD_2 , we shall focus for now on this particular step. This will then straightforwardly lead to new realisations for WA_n and WD_n .

WD_2 is generated by a current $f^{(2)}$ of spin 2 given in (2.5) and an energy-momentum tensor $T^{(2)} = W_2^{(2)}$ with central charge $2(1 + 6\alpha_0^2)$. Since $f^{(2)}$ turns out to be a spin-2 primary field with respect to $T^{(2)}$, one can define two commuting spin-2 currents

$$T_{\pm} = \frac{1}{2}(T^{(2)} \pm f^{(2)}) \quad (2.7)$$

which both generate a Virasoro algebra with the same central charge

$$c_{\pm} = 1 + 6\alpha_0^2. \quad (2.8)$$

It now follows from the reduction procedure (2.6) that the currents of $WA_3 \cong WD_3$ can be written as

$$\begin{aligned} T &= T_+ + T_- - \frac{1}{2}(\partial\varphi_3)^2 + 2\alpha_0\partial^2\varphi_3, \\ W &= \alpha_0\partial(T_+ - T_-) - \partial\varphi_3(T_+ - T_-), \\ U &= -(T_+ - T_-)^2 + 2(\partial\varphi_3)^2(T_+ + T_-) - 4\alpha_0\partial^2\varphi_3(T_+ + T_-) - 2\alpha_0\partial\varphi_3\partial(T_+ + T_-) \\ &\quad - \alpha_0^2\partial^2(T_+ + T_-) + \frac{1}{2}(1 + 6\alpha_0^2)\partial^3\varphi_3\partial\varphi_3 - \frac{2}{3}\alpha_0(1 + 6\alpha_0^2)\partial^4\varphi_3. \end{aligned} \quad (2.9)$$

Here $W = f^{(3)}$ is the spin-3 current given in (2.5), which is primary, $T = W_2^{(3)}$ is the energy-momentum tensor and U is the spin-4 current which appears in the second order pole of the operator-product expansion $W(z)W(w)$. (Note that U is given in a different basis from the spin-4 current $W_4^{(3)}$ defined in [8]; they differ by a term proportional to $\partial^2 T$.) The term $(T_+ - T_-)^2$ in U is normal ordered with respect to the modes of T_+ and T_- . One may check explicitly that these currents generate the $WA_3 \cong WD_3$ algebra with central charge

$$c = 3(1 + 20\alpha_0^2). \quad (2.10)$$

The realisation (2.9) is remarkable in the sense that T_+ and T_- may be chosen to be *arbitrary* commuting energy-momentum tensors (with equal central charges given by (2.8)). φ_3 is a free scalar which commutes with T_+ and T_- . Since these energy-momentum tensors appear in the combinations $T_+ + T_-$ as well as $T_+ - T_-$, equation (2.9) thus gives a realisation of $WA_3 \cong WD_3$ in terms of two independent energy-momentum tensors together with one free scalar field. As explained earlier, this immediately yields realisations of WA_n and WD_n in terms of two arbitrary (commuting) energy-momentum tensors $T_{\pm}$, with the same central charge (2.8), together with $(n - 2)$ free scalar fields.

These new realisations (which are at the quantum level) imply in particular the existence of new realisations of WA_n and WD_n at the classical level, *i.e.* on the level of Poisson brackets. It has been shown in [9] that Poisson-bracket realisations of WA_2 are related to Jordan algebras of cubic norm. This was generalized to a connection between Poisson-bracket realisations of WA_n and Jordan algebras of norm $\nu = n + 1$ in [10], leading to a new Poisson-bracket realisation of WA_n , which is the classical limit of the new (quantum) realisation presented in this paper.

3. Applications to String Theory

We shall now use these realisations to construct new string theories with a WA_n or WD_n symmetry. We shall prove that these W string theories are related to Virasoro minimal models. In the case of realisations with only *one* arbitrary energy-momentum tensor, it was shown in [5,6] that the corresponding W -string theories are related to the $(h, h + 1)$ unitary Virasoro minimal models, where h is the dual coxeter number of the underlying Lie algebra A_n or D_n . For the realisations with *two* arbitrary energy-momentum tensors, the corresponding W -string theories turn out to be related to two copies of the $(h, h + 1)$ unitary Virasoro minimal model. We shall treat the $WA_3 \cong WD_3$ string in some detail; the generalisation to $n \geq 4$ is then straightforward.

The critical central charge for the $WA_3 \cong WD_3$ string is $c^* = 246$. This implies that the critical value of the background-charge parameter in the Miura transformation (2.5) is, using (2.10), given by $(\alpha_0^*)^2 = 81/20$. The central charges of the energy-momentum tensors $T_{\pm}$ is then given by $c_{\pm}^* = 26 - \frac{7}{10}$. Here 26 is the critical central charge of the ordinary Virasoro string and $\frac{7}{10}$ is the central charge of the $(4, 5)$ unitary

Virasoro minimal model. This connection is strengthened by the fact that the physical spectrum of the $WA_3 \cong WD_3$ string is related to this minimal model, as we now show.

Physical states of the $WA_3 \cong WD_3$ string are defined by the conditions that they are annihilated by the positive Laurent modes of the generating currents T , W and U and that they are eigenstates of the zero Laurent modes with specific eigenvalues, *viz.* the intercepts ω_2 , ω_3 and ω_4 respectively. In principle these intercepts should be determined by a BRST analysis. A simpler method to determine them is to assume the existence of a particular “tachyonic” physical state, the “cosmological solution” [2]. It has been proven for $WA_3 \cong WD_3$, and argued for WA_n with $n \geq 4$, in [5] that the string theory based on a realisation with only one arbitrary energy-momentum tensor is unitary if and only if this cosmological solution is contained in its physical spectrum. The intercepts can thus be read off from the action of the currents on this particular state, and are given by

$$\omega_2 = 10, \quad \omega_3 = 0 \quad \text{and} \quad \omega_4 = -\frac{2607}{5}. \quad (3.2)$$

Since these intercepts depend on the basis chosen to describe the algebra, some of them differ from the ones presented in [5]. One can check, however, that the intercept values (3.2) are consistent with the ones given in [5]. In particular the spin-3 intercept in (3.2) is zero, in accordance with the fact that a spin-3 primary current always has zero intercept for any WA_n string [5].

Having obtained the intercept values, we are now in a position to construct the string theory based on this realisation. Let us consider the energy-momentum tensors $T_{\pm}$ to be realised by $D_{\pm}$ free scalars $\vec{X}_{\pm}$:

$$T_{\pm} = -\frac{1}{2}(\partial \vec{X}_{\pm})^2 + \vec{Q}_{\pm} \cdot \partial^2 \vec{X}_{\pm}, \quad (3.3)$$

where the background-charge vectors $\vec{Q}_{\pm}$ are chosen so that (2.8) is satisfied, *i.e.*

$$(\vec{Q}_{\pm})^2 = \frac{1}{12} \left(\frac{253}{10} - D_{\pm} \right). \quad (3.4)$$

For reasons we shall explain presently, all physical operators are of the form

$$V = R(\vec{X}_+, \vec{X}_-) e^{\vec{\beta}_+ \cdot \vec{X}_+ + \vec{\beta}_- \cdot \vec{X}_- + \beta_3 \varphi_3}, \quad (3.5)$$

where $R(\vec{X}_+, \vec{X}_-)$ is a differential polynomial in $\vec{X}_+$ and $\vec{X}_-$. Note in particular that physical states have no excitations in the frozen φ_3 direction. Clearly any $R(\vec{X}_+, \vec{X}_-)$ can be expressed as

$$R(\vec{X}_+, \vec{X}_-) = \sum_{i=1}^k R_+^i(\vec{X}_+) R_-^i(\vec{X}_-), \quad (3.6)$$

where k is some integer. Each differential polynomial $R_{\pm}^i(\vec{X}_{\pm})$ has level number $m_{\pm}^i$, defined by $(T_{\pm})_0 R_{\pm}^i = m_{\pm}^i R_{\pm}^i$. In order for (3.5) to satisfy the intercept conditions (3.2), one can show from (2.9) that all R_+^i must have the same level number m_+ , and all R_-^i must have the same level number m_- . The physical-state conditions then imply that (3.5) must satisfy

$$T_{\pm}(z) V(w) \sim \frac{\Delta_{\pm} V(w)}{(z-w)^2} + \frac{\partial_{\pm} V(w)}{z-w}, \quad (3.7)$$

where $\Delta_{\pm}$ are the effective intercepts for $T_{\pm}$, and ∂_+ and ∂_- stand for derivatives ∂ acting only on X_+ terms and X_- terms respectively. Note that although V is not primary under T_+ or T_- separately, it is annihilated by all positive Laurent modes of $T_{\pm}$. In fact, one can show that every physical operator V can be written as a sum of factorised products of the form

$$V_+ V_- e^{\beta_3 \varphi_3}, \quad (3.8)$$

where V_+ is built only from $\vec{X}_+$ and is primary under T_+ with dimension Δ_+ whilst V_- is built only from $\vec{X}_-$ and is primary under T_- with dimension Δ_- . Thus the theory has cloven itself into two independent Virasoro-like string theories with energy-momentum tensors $T_\pm$ having intercepts $\Delta_\pm$, and physical operators $V_\pm$.

Introducing the shifted momentum γ_3 by $\beta_3 = \alpha_0^*(2 + \gamma_3/9)$, the intercept conditions for the $WA_3 \cong WD_3$ string then give

$$(T)_0: \quad 40(\Delta_+ + \Delta_-) - \gamma_3^2 - 76 = 0, \quad (3.9a)$$

$$(W)_0: \quad \gamma_3(\Delta_+ - \Delta_-) = 0, \quad (3.9b)$$

$$(U)_0: \quad 200(\Delta_+ - \Delta_-)^2 - 20(\Delta_+ + \Delta_-)(\gamma_3^2 - 587) - 11(2028 + 23\gamma_3^2) = 0. \quad (3.9c)$$

There are six solutions to these equations, namely

Δ_+	Δ_-	γ_3
9/10	1	0
1	9/10	0
77/80	77/80	± 1
1	1	± 2

(3.10)

Thus we see that the $\Delta_\pm$ values can all be written in the form $1 - L_0^{\min}$, where $L_0^{\min}$ takes its values from the diagonal entries $\Delta_{(r,r)}$ of the Kac table of the $(4, 5)$ minimal model.

It is now straightforward to demonstrate the unitarity of the theory since it is effectively described by two independent Virasoro-like strings, each with central charge $26 - \frac{7}{10}$ and intercept values given in (3.10). Since these lie within the unitarity bounds for the $c = 26 - \frac{7}{10}$ string, the absence of ghosts is manifest.

In the discussions so far we have only considered states that have no excitations in the frozen direction φ_3 . Following [5], we can show that the momentum-conservation law in the φ_3 direction, which must be satisfied in order to have non-vanishing two-point functions, requires that the physical operators should come in conjugate pairs with equal and opposite shifted momenta γ_3 . For the physical operators we have considered so far, this is indeed the case, as one can see in (3.10). Explicit computations for level 1 states, combined with evidence found previously for other W strings [5,6], indicate that this conjugation property is never satisfied for states with excitations in the frozen direction. Thus such states have vanishing two-point functions and therefore have zero norm. Consequently, the only physical states are indeed given by sums of terms of the form (3.8).

4. Conclusions

In this paper we have described new realisations for the WA_n and WD_n algebras for $n \geq 3$. The currents are realised in terms of two arbitrary energy-momentum tensors together with $(n - 2)$ free scalars. We have discussed in detail the example of $WA_3 \cong WD_3$, and given the explicit form of the currents. We have used this realisation to construct a new W string. The essential new feature of this string theory is that it is effectively reduced to two copies of Virasoro-like strings with the same central charge $26 - \frac{7}{10}$. These strings are independent except that their intercepts are correlated, as can be seen in (3.10).

This effective theory is very similar to the theory that one would obtain by starting from an action that was the sum of two independent Virasoro-string actions. Thus it describes two “non-communicating” worlds, each with its own spacetime.

Although we have concentrated for simplicity on the $n = 3$ case, the generalisation to WA_n and WD_n for arbitrary n is completely straightforward, using the reduction procedures explained in section 2. The

intercepts for the two effective Virasoro-string theories can easily be found from the action of the Weyl group of A_n or D_n on the cosmological solution in the n -scalar realisation, using the same techniques that were developed in [5,6]. These effective intercepts are again related to the diagonal entries of the Kac tables for the $(h, h+1)$ unitary Virasoro minimal models, where h is the dual Coxeter number of the underlying Lie algebra.

ACKNOWLEDGMENTS

Stany Schrans is much obliged to the Center for Theoretical Physics, Texas A&M University, for hospitality, and to the Belgian National Fund for Scientific Research for a travel grant.

REFERENCES

- [1] C.N. Pope, L.J. Romans and K.S. Stelle, *Phys. Lett.* **268B** (1991) 167;
Phys. Lett. **269B** (1991) 287.
- [2] S.R. Das, A. Dhar and S.K. Rama, *Mod. Phys. Lett.* **A6** (1991) 3055;
“Physical states and scaling properties of W gravities and W strings,” TIFR/TH/91-20.
- [3] C.N. Pope, L.J. Romans, E. Sezgin and K.S. Stelle, *Phys. Lett.* **274B** (1992) 298.
- [4] S.K. Rama, *Mod. Phys. Lett.* **A6** (1991) 3531.
- [5] H. Lu, C.N. Pope, S. Schrans and K.W. Xu, “The Complete Spectrum of the W_N String,” preprint CTP TAMU-5/92, KUL-TF-92/1.
- [6] H. Lu, C.N. Pope, S. Schrans and X.J. Wang, “On Sibling and Exceptional W Strings,” preprint CTP TAMU-10/92, KUL-TF-92/8.
- [7] V.A. Fateev and S. Lukyanov, *Int. J. Mod. Phys.* **A3** (1988) 507.
- [8] S.L. Lukyanov and V.A. Fateev, *Sov. Scient. Rev.* **A15** (1990) 1;
Sov. J. Nucl. Phys. **49** (1989) 925.
- [9] L.J. Romans, *Nucl. Phys.* **B352** (1991) 829.
- [10] C.M. Hull, *Nucl. Phys.* **B353** (1991) 707.